



## **Seguimento de indivíduos com dispositivos GPS**



0 5 10 20 Kilometers

**Larus michahellis (incubation)**


- Lmic\_3510    Lmic\_217    Lmic\_246    Lmic\_15
- Lmic\_3509    Lmic\_210    Lmic\_138    Lmic\_08
- Lmic\_3505    Lmic\_261    Lmic\_126    Lmic\_05
- Lmic\_3504



0 12.5 25 50 Kilometers

**Larus michahellis (post-breeding)**


- 
 Lmic\_8
  Lmic\_6
  Lmic\_4
  Lmic\_2
- 
 Lmic\_7
  Lmic\_5
  Lmic\_3



0 250 500 1,000 Kilometers

**Calonectris borealis (pre-laying)**

- |                 |                 |                 |                 |                 |
|-----------------|-----------------|-----------------|-----------------|-----------------|
| — Cbor_pl_0     | — Cbor_pl_39109 | — Cbor_pl_59751 | — Cbor_pl_64392 | — Cbor_pl_76506 |
| — Cbor_pl_22523 | — Cbor_pl_39114 | — Cbor_pl_64245 | — Cbor_pl_64393 | — Cbor_pl_76516 |
| — Cbor_pl_29047 | — Cbor_pl_55488 | — Cbor_pl_64291 | — Cbor_pl_64397 | — Cbor_pl_78298 |
| — Cbor_pl_55495 | — Cbor_pl_64336 | — Cbor_pl_73001 | — Cbor_pl_78352 |                 |



*Calonectris borealis* (chick rearing)

0 50 100 200 Kilometers

- |                 |                 |                 |                 |                 |
|-----------------|-----------------|-----------------|-----------------|-----------------|
| — Cbor_ch_0     | — Cbor_ch_55488 | — Cbor_ch_64361 | — Cbor_ch_67783 | — Cbor_ch_76516 |
| — Cbor_ch_7656  | — Cbor_ch_59751 | — Cbor_ch_64393 | — Cbor_ch_73001 | — Cbor_ch_76603 |
| — Cbor_ch_39109 | — Cbor_ch_62826 | — Cbor_ch_64397 | — Cbor_ch_73262 | — Cbor_ch_78292 |
| — Cbor_ch_39114 | — Cbor_ch_62839 | — Cbor_ch_67654 | — Cbor_ch_73269 | — Cbor_ch_78298 |
| — Cbor_ch_44539 | — Cbor_ch_64291 | — Cbor_ch_67668 | — Cbor_ch_73290 | — Cbor_ch_78615 |
| — Cbor_ch_54006 |                 |                 |                 |                 |